arXiv:1601.01421v2 [cs.IT] 22 Jan 2016

Repeated-root constacyclic codes of length $3lp^s$ and their dual codes *

Li Liu, Lanqiang Li, Xiaoshan Kai, Shixin Zhu

School of Mathematics, Hefei University of Technology, Hefei 230009, Anhui, P.R.China

Abstract: Let $p \neq 3$ be any prime and $l \neq 3$ be any odd prime with $\gcd(p, l) = 1$. The multiplicative group $F_q^* = \langle \xi \rangle$ can be decomposed into mutually disjoint union of $\gcd(q-1, 3lp^s)$ cosets over the subgroup $\langle \xi^{3lp^s} \rangle$, where ξ is a primitive $(q-1)$th root of unity. We classify all repeated-root constacyclic codes of length $3lp^s$ over the finite field F_q into some equivalence classes by this decomposition, where $q = p^m$, s and m are positive integers. According to these equivalence classes, we explicitly determine the generator polynomials of all repeated-root constacyclic codes of length $3lp^s$ over F_q and their dual codes. Self-dual cyclic codes of length $3lp^s$ over F_q exist only when $p = 2$. And we give all self-dual cyclic codes of length $3 \cdot 2^s l$ over F_{2^m} and their enumeration.

Keywords: Repeated-root constacyclic codes, Cyclic codes, Dual codes, Generator polynomial

1 Introduction

Constacyclic codes over finite fields play a very important role in the theory of error-correcting codes. More importantly, constacyclic codes have practical applications. As these codes have rich algebraic structures, they can be efficiently encoded and decoded using shift registers. They also have very good error-correcting properties. All of those explain their preferred role in engineering.

Repeated-root cyclic codes were first investigated in the 1990s by Castagnoli in [1] and Van Lint in [2], where it was proved that repeated-root cyclic codes have a concatenated construction, and are asymptotically bad. However, it is well known that there still exist a few optimal such codes (see $[12-14]$), which encourages many scholars to study the class of codes. For example, Dinh determined the generator polynomials of all constacyclic codes and their dual codes over F_q of length $2p^s, 3p^s$ and $6p^s$ in $[3-5]$. Since then, these results have been extended to more general code lengths. In 2012, Bakshi and Raka gave the generator polynomials of all constacyclic codes of length $2^t p^s$ over F_q in [6], where q is a power of an odd prime p. In 2014, Chen et.al studied all constacyclic codes of length lp^s over F_q in [7], where l is a prime different from p. In [7], all constacyclic codes of length lp^s over F_q and their dual codes were obtained, and all self-dual and all linear complementary dual constacyclic codes were given. Recently, in [8], Sharma explicitly determined the generator polynomials of all repeated-root constacyclic codes of length $l^t p^s$ over F_{p^m} and their dual codes. Further, they listed all self-dual cyclic and negacyclic codes and also determined all self-orthogonal cyclic and negacyclic codes of length $l^t p^s$ over F_{p^m}. What's more, Chen et.al studied all constacyclic codes of length $2l^m p^s$ over F_q of characteristic p in [9], and they gave the characterization and enumeration of all linear complementary dual and self-dual constacyclic codes of length $2l^m p^s$ over F_q. In the conclusion of their paper, they proposed to study all constacyclic codes of length $kl^m p^s$ over F_q, where p is the characteristic of F_q, l is an odd prime different from p, and k is a prime different from l and p. However, this is not an easy work.

* E-mail addresses: liuli-1128@163.com(Li Liu), lilanqiang716@126.com(Lanqiang Li).
This research is supported by the National Natural Science Foundation of China (No.61572168,No.11401154) and the Anhui provincial Natural Science Foundation under Grant (No.158085MA13).

In this paper, we study all constacyclic codes of length $3lp^s$ over F_q, where $p \neq 3$ is any prime and $l \neq 3$ is any odd prime with $\gcd(p, l) = 1$. The article is organized as follows. In section 3, we decompose the multiplicative cyclic group $F_q^* = \langle \xi \rangle$ into mutually disjoint union of cosets of $\langle \xi^{3lp^s} \rangle$, which are one-to-one correspondence to the equivalence classes of all constacyclic. Based on this decomposition, Section 4 explicitly determines the generator polynomials of all $\lambda-$constacyclic codes of length $3lp^s$ over F_q and their dual codes, where λ is any none-zero element of F_q and $q = p^m$ is a power of prime. As an application, we give all self-dual cyclic codes of length $3 \cdot 2^s l$ over F_{2^m} and their enumeration in Section 5.

2 Preliminaries

Let F_q be the finite field of order $q = p^m$, where $p \neq 3$ is a prime and the characteristic of the field, and m is a positive integer. Let $F_q^* = \langle \xi \rangle$ be the multiplicative cyclic group of none-zero elements of F_q, where ξ is a primitive $(q-1)$th root of unity.

For any element $\lambda \in F_q^*$, $\lambda-$constacyclic codes of length n over F_q are regarded as the ideals $\langle g(x) \rangle$ of the quotient ring $F_q[x]/(x^n - \lambda)$, where $g(x)|x^n - \lambda$. Further, the definition of the dual code of code C is as follows:

$$C^{\perp} = \{x \in F_q^n | x \cdot y = 0, \forall y \in C\},$$

where $x \cdot y$ denotes the Euclidean inner product of x and y in F_q^n. The code C is called to be a self-orthogonal code if $C \subseteq C^{\perp}$ and a self-dual code if $C = C^{\perp}$. Let C be a $\lambda-$constacyclic code of length n over F_q generated by a polynomial $g(x)$, i.e., $C = \langle g(x) \rangle$. As $g(x)|x^n - \lambda$, then there must be a polynomial $h(x) \in F_q[x]$ such that $h(x) = \frac{x^n - \lambda}{g(x)}$. It is clear that $h(x)$ is also monic if $g(x)$ is monic. The polynomial $h(x)$ is called the parity check polynomial of code C. It is well known that the dual code $C^{\perp}$ is generated by $h(x)^*$, where $h(x)^*$ is the reciprocal polynomial of $h(x)$. For any $f(x) \in F_q[x]$, the reciprocal polynomial of $f(x)$ is defined as $f(x)^* = f(0)^{-1} x^{\deg(f(x))} f(\frac{1}{x})$. Obviously, $(f_1 f_2)^* = f_1^* f_2^*$, and $(f_1^*)^* = f_1$, for any polynomials $f_1(x), f_2(x) \in F_q[x]$.

Let n be any positive integer. For any integer s, $0 \leq s \leq n-1$, we have the definition of $q-$cyclotomic coset of s modulo n is as follows:

$$C_s = \{s, sq, ..., sq^{n_s - 1}\},$$

where n_s is the least positive integer such that $sq^{n_s} \equiv s (\text{mod } n)$. Then it is easy to see that n_s is equal to the multiplicative order of q modulo $\frac{n}{\gcd(s,n)}$. If α denotes a primitive nth root of unity in some extension field of F_q, then the polynomial $M_s(x) = \prod_{i \in C_s} (x - \alpha^i)$ is the minimal polynomial of α^s over F_q, and

$$x^n - 1 = \prod M_s(x)$$

gives the factorization of $(x^n - 1)$ into irreducible factors over F_q, where s runs over a complete set of representatives from distinct $q-$cyclotomic coset modulo n.

Obviously, when $n = l$, where $l \neq 3$ is an odd prime with $\gcd(l, p) = 1$, we get that all the distinct $q-$cyclotomic cosets modulo l are $C_0 = \{0\}$ and $C_k = \{g^k, g^k q, ..., g^k q^{n_k - 1}\}$, for any integer k, $1 \leq k \leq e = \frac{\phi(l)}{f}$, by [15, Theorem 1], where g is a fixed generator of the cyclic group Z_l^*, $f = \text{ord}_l(q)$ is the multiplicative order of q in Z_l^*, and ϕ is Euler's phi-function. Therefore, the irreducible factorization of $x^l - 1$ over F_q is given by

$$x^l - 1 = M_0(x) M_1(x) M_2(x) ... M_e(x),$$

where $M_i(x) = \prod_{j \in C_i}(x - \eta^i)$ with η being a primitive lth root of unity.

In addition, we determine all the distinct q^3-cyclotomic cosets modulo l, which is needed to prove our main results. There exist two subcases. When $\gcd(f, 3) = 1$, it is easy to prove that $f = \mathrm{ord}_l(q) = \mathrm{ord}_l(q^3)$, then $\langle q \rangle = \langle q^3 \rangle$ in Z_l^*. According to the definition of q^3-cyclotomic coset modulo l, we have that C_0 and C_k, $1 \leq k \leq e = \frac{\phi(l)}{f}$, are also all of the distinct q^3-cyclotomic cosets modulo l. When $\gcd(f, 3) = 3$, we prove that $\mathrm{ord}_l(q^3) = \frac{f}{3}$. It is easy to verify that $A_0 = \{0\}$,

$$A_k = \{g^k, g^k q^3, ..., g^k q^{3(\frac{f}{3}-1)}\},$$

$$A_{kq} = \{g^k q, g^k q q^3, ..., g^k q q^{3(\frac{f}{3}-1)}\},$$

$$A_{kq^2} = \{g^k q^2, g^k q^2 q^3, ..., g^k q^2 q^{3(\frac{f}{3}-1)}\},$$

consist of all the distinct q^3-cyclotomic cosets modulo l, where $1 \leq k \leq e$. Then we have the irreducible factorization of $x^l - 1$ in $F_{q^3}[x]$ as follows:

$$x^l - 1 = A_0(x)A_1(x)A_q(x)A_{q^2}(x)A_2(x)A_{2q}(x)A_{2q^2}(x)...A_e(x)A_{eq}(x)A_{eq^2}(x),$$

where $A_0(x) = (x - 1)$, $A_k(x) = \prod_{s \in A_k}(x - \eta^s)$, $A_{kq}(x) = \prod_{t \in A_{kq}}(x - \eta^t)$ and $A_{kq^2}(x) = \prod_{j \in A_{kq^2}}(x - \eta^j)$, $1 \leq k \leq e$.

We also give all the distinct $q-$cyclotomic cosets modulo $3l$. As $\gcd(q, 3) = 1$, we have $q^{\phi(3)} \equiv 1 (\mathrm{mod}\ 3)$ by Euler's Theorem, i.e., $q^2 \equiv 1 (\mathrm{mod}\ 3)$. Then, it is simple to verify that

$$\mathrm{ord}_{3l}(q) = \begin{cases} f, & q \equiv 1 (\mathrm{mod}\ 3); \\ f, & q \equiv 2 (\mathrm{mod}\ 3) \text{ with } f \text{ even}; \\ 2f, & q \equiv 2 (\mathrm{mod}\ 3) \text{ with } f \text{ odd}. \end{cases}$$

From [16, Chapter 8], there exists a primitive root r modulo l such that $\gcd(\frac{r^{l-1}-1}{l}, l) = 1$. Assume that $g = r + (1 - r)l^2$, we have $g^{l-1} - 1 \equiv (r + (1 - r)l^2)^{l-1} - 1 \equiv r^{l-1} - 1 (\mathrm{mod}\ l^2)$. Therefore, $\gcd(\frac{g^{l-1}-1}{l}, l) = \gcd(\frac{r^{l-1}-1}{l}, l) = 1$. It's clear that g is a primitive root modulo l^t, $t \geq 1$, such that $g \equiv 1 (\mathrm{mod}\ 3)$.

We give all the distinct $q-$cyclotomic cosets modulo $3l$ by the following lemma.

Lemma 2.1. (I) If $q \equiv 1 (\mathrm{mod}\ 3)$, then we have that all the distinct $q-$cyclotomic cosets modulo $3l$ are given by

$$B_0 = \{0\}, B_l = \{l\}, B_{-l} = \{-l\},$$

$$B_{ag^k} = \{ag^k, ag^k q, ..., ag^k q^{f-1}\},$$

for $a \in R = \{1, -1, 3\}$ and $0 \leq k \leq e - 1$.
(II) If $q \equiv 2 (\mathrm{mod}\ 3)$ and f is even, we have that all the distinct $q-$cyclotomic cosets modulo $3l$ are given by $B_0 = \{0\}, B_l = \{l, lq\}$,

$$B_{g^{k'}} = \{g^{k'}, g^{k'} q, ..., g^{k'} q^{f-1}\}, \text{ for } 0 \leq k' \leq 2e - 1,$$

$$B_{3g^k} = \{3g^k, 3g^k q, ..., 3g^k q^{f-1}\}, \text{ for } 0 \leq k \leq e - 1.$$

(III) If $q \equiv 2 (\mathrm{mod}\ 3)$ and f is odd, we have that all the distinct $q-$cyclotomic cosets modulo $3l$ are given by

$$B_0 = \{0\}, B_l = \{l, lq\},$$

$$B_{g^k} = \{g^k, g^k q, ..., g^k q^{2f-1}\},$$
$$B_{3g^k} = \{3g^k, 3g^k q, ..., 3g^k q^{f-1}\},$$

for $0 \leq k \leq e-1$.

Proof. (I) Firstly, we prove that the cyclotomic cosets $B_{ag^k}, 0 \leq k \leq e-1$, are distinct. If there exist some k_1, k_2, $0 \leq k_1, k_2 \leq e-1$, such that $B_{ag^{k_1}} = B_{ag^{k_2}}$, then we have

$$a_1 g^{k_1} \equiv a_2 g^{k_2} q^j (\text{mod } 3l),$$

for some integer j, where $a_1, a_2 \in R = \{1, -1, 3\}$. Therefore, we get

$$\gcd(a_1 g^{k_1}, 3l) = \gcd(a_2 g^{k_2} q^j, 3l) = \gcd(a_2 g^{k_2}, 3l),$$

as $\gcd(q, 3l) = 1$. From this, we can deduce $a_1 = a_2$ or $a_1 = -a_2 = \pm 1$.

If $a_1 = -a_2 = \pm 1$, then

$$-g^{k_1} \equiv g^{k_2} q^j (\text{mod } 3l), \quad \text{i.e.,} \quad -1 \equiv g^{k_1 - k_2} q^j (\text{mod } 3l),$$

for some integer j. Due to $g \equiv 1(\text{mod } 3)$ and $q \equiv 1(\text{mod } 3)$, we deduce $-1 \equiv 1(\text{mod } 3)$. This is a contradiction.

If $a_1 = a_2$, assume that $a_1 = a_2 = a$, then we have

$$ag^{k_1} \equiv ag^{k_2} q^j (\text{mod } 3l), \quad \text{i.e.,} \quad g^{k_1 - k_2} \equiv q^j (\text{mod } l),$$

for some integer j. Further, we have $g^{(k_1-k_2)f} \equiv q^{jf} \equiv 1(\text{mod } l)$. As g is a primitive root modulo l, we get $\phi(l) | (k_1 - k_2)f$, i.e., $\text{e} = \frac{\phi(l)}{f} | k_1 - k_2$. Since $0 \leq k_1, k_2 \leq e-1$, we must have $k_1 = k_2$. Secondly, we get

$$\begin{aligned} |B_0| + |B_l| + |B_{-l}| + \sum_{a \in R} \sum_{k=0}^{e-1} |B_{ag^k}| &= 3 + \sum_{a \in R} \sum_{k=0}^{e-1} f \\ &= 3 + \sum_{a \in R} ef \\ &= 3 + 3\phi(l) \\ &= 3l. \end{aligned}$$

So, the conclusion (I) holds. The conclusions (II) and (III) are also established in a similar way. □

Assume that $B_o(x), B_l(x), B_{-l}(x)$ and $B_{ag^k}(x)$ are the minimal polynomials of the corresponding cosets B_o, B_l, B_{-l} and B_{ag^k}. From the above lemma, we get the following theorem immediately.

Theory 2.2. The irreducible factorization of $x^{3l} - 1$ over F_q is given as follows:
(I) If $q \equiv 1(\text{mod } 3)$, then

$$x^{3l} - 1 = B_0(x) B_l(x) B_{-l}(x) \prod_{a \in R} \prod_{k=0}^{e-1} B_{ag^k}(x),$$

where $a \in R = \{1, -1, 3\}$ and $0 \leq k \leq e-1$.
(II) If $q \equiv 2(\text{mod } 3)$ and f is even, then

$$x^{3l} - 1 = B_0(x)B_l(x) \prod_{k'=0}^{2e-1} B_{g^{k'}}(x) \prod_{k=0}^{e-1} B_{3g^k}(x),$$

where $0 \leq k \leq e-1, 0 \leq k^{'} \leq 2e-1$.
(III) If $q \equiv 2(\text{mod } 3)$ and f is odd, then

$$x^{3l} - 1 = B_0(x)B_l(x) \prod_{k=0}^{e-1} B_{g^k}(x)B_{3g^k}(x),$$

where $0 \leq k \leq e-1$.

The next two lemmas give the necessary and sufficient conditions for judging the reducibility of binomials and trinomial, which were given by Wan in [17].

Lemma 2.3. Suppose that $n \geq 2$, Let $k = \text{ord}(a)$ be the multiplicative order of a, for any $a \in F_q^*$. Then, the binomial $x^n - a$ is irreducible over F_q if and only if
(i) Every prime divisor of n divides k, but not $\frac{(q-1)}{k}$;
(ii) If $4|n$, then $4|(q-1)$.

Lemma 2.4. Let t be a positive integer, and $H(x) \in F_q[x]$ be irreducible over F_q with $\deg(H(x)) = n$, x does not divide $H(x)$, and e denotes the order of any root of $H(x)$. Then $H(x^t)$ is irreducible over F_q if and only if
(i) Each prime divisor of t divides e;
(ii) $\gcd(t, \frac{q^n-1}{e}) = 1$;
(iii) If $4|t$, then $4|(q^n-1)$.

3 A classification of constacyclic codes of length $3lp^s$

Let ξ be a primitive $(q-1)$th root of unity, and $F_q^* = \langle \xi \rangle$ be a cyclic group of order $(q-1)$ as before. It is easy to verify that $\langle \xi^{3lp^s} \rangle = \langle \xi^{3l} \rangle = \langle \xi^d \rangle$ and the index $|F_q^* : \langle \xi^{3lp^s} \rangle| = d$, where $d = \gcd(q-1, 3lp^s)$. Thus, the multiplicative cyclic group F_q^* can be decomposed into mutually disjoint union of cosets over the subgroup $\langle \xi^{3lp^s} \rangle$ as follows:

Lemma 3.1. $F_q^* = \langle \xi \rangle = \langle \xi^d \rangle \bigcup \xi^{p^s} \langle \xi^d \rangle \bigcup ... \bigcup \xi^{p^s(d-1)} \langle \xi^d \rangle$, where $d = \gcd(q-1, 3lp^s)$.

According to the properties of the coset, we obtain the following lemma immediately.

Lemma 3.2. For any two none-zero elements λ and μ of F_q, there exists some integer j, $0 \leq j \leq d-1$ such that $\lambda, \mu \in \xi^{jp^s} \langle \xi^d \rangle$ if and only if $\lambda^{-1}\mu \in \langle \xi^d \rangle$, where $d = \gcd(q-1, 3lp^s)$.

If λ and μ are in the same coset, we build a one-to-one correspondence between $\lambda-$constacyclic codes and $\mu-$constacyclic codes of length $3lp^s$ over F_q as following theorem, which shows that $\lambda-$constacyclic codes and $\mu-$constacyclic codes are equivalent.

Theorem 3.3. Let $\lambda, \mu \in F_q^*$. Then there exists some integer $a \in F_q^*$ such that

$$\varphi : F_q[x]/(x^{3lp^s} - \mu) \to F_q[x]/(x^{3lp^s} - \lambda),$$
$$f(x) \mapsto f(ax),$$

is an isomorphism if and only if $\lambda, \mu \in \xi^{jp^s}\langle \xi^d \rangle$, where $0 \le j \le d-1$.

Proof. ($\Rightarrow$) If φ is an isomorphism, then we have

$$\mu = \varphi(\mu) = \varphi(x^{3lp^s}) = (\varphi(x))^{3lp^s} = (ax)^{3lp^s} = a^{3lp^s} x^{3lp^s} = a^{3lp^s}\lambda,$$

i.e., $\lambda^{-1}\mu = a^{3lp^s}$. As $a = \xi^k \in F_q^*$, for some positive integer k, then $\lambda^{-1}\mu = \xi^{k\cdot 3lp^s} \in \langle \xi^d \rangle$. By Lemma 3.2, we get that there exists j, $0 \le j \le d-1$, such that $\lambda, \mu \in \xi^{jp^s}\langle \xi^d \rangle$.

($\Leftarrow$) If there exists j, $0 \le j \le d-1$, such that $\lambda, \mu \in \xi^{jp^s}\langle \xi^d \rangle$, then we have $\lambda^{-1}\mu \in \langle \xi^d \rangle = \langle \xi^{3lp^s} \rangle$, by Lemma 3.2 again. Thus, $\lambda^{-1}\mu = \xi^{k\cdot 3lp^s}$, for some integer k. Set $a = \xi^k$, then $\lambda a^{3lp^s} = \mu$. Further, it is easy to prove that the following map is an isomorphism:

$$\varphi : F_q[x]/(x^{3lp^s} - \mu) \to F_q[x]/(x^{3lp^s} - \lambda),$$
$$f(x) \mapsto f(ax).$$

□

From Theorem 3.3, we get the following obvious corollaries.

Corollary 3.4. For any two elements λ and μ of F_q^*, a $\lambda-$constacyclic code is equivalent to an $\mu-$constacyclic code if and only if there exists j, $0 \le j \le d-1$, such that $\lambda, \mu \in \xi^{jp^s}\langle \xi^d \rangle$. Further, a $\lambda-$constacyclic code and an $\mu-$constacyclic code are both equivalent to a $\xi^{jp^s}-$constacyclic code.

Corollary 3.5. Let λ be any element of F_q^*, then there exists some integer j, $0 \le j \le d-1$, such that a $\lambda-$constacyclic code is equivalent to a $\xi^{jp^s}-$constacyclic code.

Obviously, the Theorem 3.3 and its two corollaries show that all constacyclic codes of length $3lp^s$ over F_q are classified into $d = \gcd(q-1, 3lp^s)$ mutually disjoint classes. It is enough to consider $\lambda-$constacyclic codes, where $\lambda = \xi^{jp^s}$, $0 \le j \le d-1$, and $d = \gcd(q-1, 3lp^s)$, if we want to determine all constacyclic codes of length $3lp^s$ over F_q. Therefore, we mainly study $\lambda-$constacyclic codes in Section 4.

4 All constacyclic codes of length $3lp^s$ over F_q

Let $f(x)$ be any polynomial of $F_q[x]$ and leading coefficient $a_n \neq 0$, we denote $\widehat{f}(x) = a_n^{-1} f(x)$. Then, $\widehat{f}(x)$ is said to be the monic polynomial of $f(x)$.

From the above discussion in the section 3, we know that the number of equivalence constacyclic classes is equal to $d = \gcd(q-1, 3lp^s)$. It is obvious that the value of d has the following four cases:
(i) $d = \gcd(q-1, 3lp^s) = 1$.
(ii) $d = \gcd(q-1, 3lp^s) = 3$.
(iii) $d = \gcd(q-1, 3lp^s) = l$.
(iv) $d = \gcd(q-1, 3lp^s) = 3l$.

4.1 All constacyclic codes of length $3lp^s$ over F_q when $d = 1$

From Lemma 3.1, we see that $F_q^* = \langle \xi \rangle$ is the decomposition of coset over the subgroup $\langle \xi^d \rangle$, when $d = \gcd(q-1, 3lp^s) = 1$. In this case, it is clear that all constacyclic codes of length $3lp^s$ over F_q are equivalent to the cyclic codes. Therefore, we have the following theorem.

Theorem 4.1. Let $d = \gcd(q-1, 3lp^s) = 1$, then $\lambda-$constacyclic codes C of length $3lp^s$ over F_q are equivalent to the cyclic codes, for any $\lambda \in F_q^*$, i.e., there exists a unique element $a \in F_q^*$ such that $a^{3lp^s}\lambda = 1$. Further, we have the map

$$\begin{aligned} &\varphi_a : F_q[x]/(x^{3lp^s} - 1) \to F_q[x]/(x^{3lp^s} - \lambda), \\ &f(x) \mapsto f(ax), \end{aligned}$$

is an isomorphism, and the irreducible factorization of $x^{3lp^s} - \lambda$ in $F_q[x]$ is given by:
(I) If f is even, then

$$x^{3lp^s} - \lambda = \widehat{B}_0(ax)^{p^s} \widehat{B}_l(ax)^{p^s} \prod_{k'=0}^{2e-1} \widehat{B}_{g^{k'}}(ax)^{p^s} \prod_{k=0}^{e-1} \widehat{B}_{3g^k}(ax)^{p^s},$$

where $0 \le k \le e-1, 0 \le k' \le 2e-1$.

Therefore, we have

$$C = \langle \widehat{B}_0(ax)^{\varepsilon} \widehat{B}_l(ax)^{\rho} \prod_{k'=0}^{2e-1} \widehat{B}_{g^{k'}}(ax)^{\tau_{k'}} \prod_{k=0}^{e-1} \widehat{B}_{3g^k}(ax)^{\upsilon_k} \rangle,$$

$$C^{\perp} = \langle \widehat{B}_0(a^{-1}x)^{p^s-\varepsilon} \widehat{B}_{-l}(a^{-1}x)^{p^s-\rho} \prod_{k'=0}^{2e-1} \widehat{B}_{-g^{k'}}(a^{-1}x)^{p^s-\tau_{k'}} \prod_{k=0}^{e-1} \widehat{B}_{-3g^k}(a^{-1}x)^{p^s-\upsilon_k} \rangle,$$

where $0 \le \varepsilon, \rho, \tau_{k'}, \upsilon_k \le p^s$, for any $k = 0, 1, 2, ..., e-1$, and $k' = 0, 1, 2, ..., 2e-1$.
(II) If f is odd, then

$$x^{3lp^s} - \lambda = \widehat{B}_0(ax)^{p^s} \widehat{B}_l(ax)^{p^s} \prod_{k=0}^{e-1} \widehat{B}_{g^k}(ax)^{p^s} \widehat{B}_{3g^k}(ax)^{p^s},$$

where $0 \le k \le e-1$.

Therefore, we have

$$C = \langle \widehat{B}_0(ax)^{\varepsilon} \widehat{B}_l(ax)^{\rho} \prod_{k=0}^{e-1} \widehat{B}_{g^k}(ax)^{\tau_k} \widehat{B}_{3g^k}(ax)^{\upsilon_k} \rangle,$$

$$C^{\perp} = \langle \widehat{B}_0(a^{-1}x)^{p^s-\varepsilon} \widehat{B}_{-l}(a^{-1}x)^{p^s-\rho} \prod_{k=0}^{e-1} \widehat{B}_{-g^k}(a^{-1}x)^{p^s-\tau_k} \widehat{B}_{-3g^k}(a^{-1}x)^{p^s-\upsilon_k} \rangle,$$

where $0 \le \varepsilon, \rho, \tau_k, \upsilon_k \le p^s$, for any $k = 0, 1, 2, ..., e-1$.

Proof. From Theorems 2.2 and 3.3, we see that this theorem is obtained immediately. □

4.2 All constacyclic codes of length $3lp^s$ over F_q when $d = 3$

In this subsection, we consider the second case, i.e., $d = \gcd(q-1, 3lp^s) = 3$. In this case, we have that $F_q^* = \langle \xi \rangle = \langle \xi^3 \rangle \bigcup \xi^{p^s} \langle \xi^3 \rangle \bigcup \xi^{2p^s} \langle \xi^3 \rangle$ is the decomposition of cosets of F_q^* over subgroup $\langle \xi^{3lp^s} \rangle$ by Lemma 3.1. Therefore, it is enough to consider the cyclic codes, $\xi^{p^s}-$constcyclic codes and $\xi^{2p^s}-$constcyclic codes if we want to determine all constacyclic codes of length $3lp^s$ over F_q, when $\gcd(q-1, 3lp^s) = 3$.

From the Section 2, we have that the irreducible factorization of $x^l - 1$ is given by $x^l - 1 = \prod_{i=0}^{e} M_i(x)$, where $M_i(x) = \prod_{j \in C_i}(x - \eta^i)$ with η is a primitive lth root of unity. According to this, we deduce the following lemma, which gives the irreducible factorization of $x^{lp^s} - \xi^j$ in $F_q[x]$, when $\gcd(q-1, l) = 1$.

Lemma 4.2. Let $\gcd(q-1, l) = 1$. Then, for any $\xi^j \in F_q^*$, there exists a unique element $b_j \in F_q^*$ such that $b_j^{lp^s} \xi^j = 1$, where $j = 0, 1, 2, ...q-1$. Thus, the irreducible factorization of $x^{lp^s} - \xi^j$ is given by

$$x^{lp^s} - \xi^j = \prod_{i=0}^{e} \widehat{M_i}(b_j x)^{p^s},$$

where $M_i(x)$ is the minimal polynomial of the $q-$cyclotomic coset C_i modulo l.

Proof. Similar to Lemma 3.1, we have the decomposition of coset of $F_q^* = \langle \xi \rangle$ over the subgroup $\langle \xi^{lp^s} \rangle$ is given by $F_q^* = \langle \xi \rangle$, when $\gcd(q-1, l) = 1$. Next, by Theorem 3.3, we have the result. □

Lemma 4.3. Let $\gcd(q-1, 3lp^s) = 3$. Then the irreducible factorization of $x^{3lp^s} - 1$ is given by

$$x^{3lp^s} - 1 = \prod_{i=0}^{e} M_i(x)^{p^s} \widehat{M_i}(b_{\frac{q-1}{3}} x)^{p^s} \widehat{M_i}(b_{\frac{2(q-1)}{3}} x)^{p^s},$$

where $M_i(x)$ is the minimal polynomial of the $q-$cyclotomic coset C_i modulo l and $b_{\frac{q-1}{3}}$, $b_{\frac{2(q-1)}{3}} \in F_q^*$.

Proof. As $\gcd(q-1, 3lp^s) = 3$, then it is clear that $\xi^{\frac{q-1}{3}} \in F_q^*$ is a primitive 3th root of unity. Hence, we have

$$x^{3lp^s} - 1 = (x^l - 1)^{p^s} (x^l - \xi^{\frac{q-1}{3}})^{p^s} (x^l - \xi^{\frac{2(q-1)}{3}})^{p^s}.$$

Further, by Lemma 4.2, we get

$$x^{3lp^s} - 1 = \prod_{i=0}^{e} M_i(x)^{p^s} \widehat{M_i}(b_{\frac{q-1}{3}} x)^{p^s} \widehat{M_i}(b_{\frac{2(q-1)}{3}} x)^{p^s},$$

where $M_i(x)$ is the minimal polynomial of the $q-$cyclotomic coset C_i modulo l and $b_{\frac{q-1}{3}}$, $b_{\frac{2(q-1)}{3}} \in F_q^*$. □

Before determining $\xi^{ip^s}-$constacyclic codes, $i = 1, 2$, we must explicitly decompose the polynomial $x^{3lp^s} - \xi^{ip^s}$ $(i = 1, 2)$, into the product of monic irreducible factors. Obviously, we only need to determine the irreducible factorization of $x^{3l} - \xi^i$ $(i = 1, 2)$. Firstly, we consider the polynomial $x^3 - \xi^i$ $(i = 1, 2)$. Since, $x^3 - \xi^i$ $(i = 1, 2)$ is irreducible in $F_q[x]$, we get that F_{q^3} is a splitting field for $x^3 - \xi^i$ $(i = 1, 2)$ over F_q. Thus, there exists $\nu_i \in F_{q^3}$ such that $\nu_i^3 = \xi^i$ $(i = 1, 2)$. Further, it is easy to get that $\nu_i, \alpha\nu_i$, and $\alpha^2\nu_i$ are all the roots of $x^3 - \xi^i$ $(i = 1, 2)$, in F_{q^3}, where α is a primitive 3th root of unity. In addition, we see that $\nu_i \in F_{q^3}$ but $\nu_i \notin F_q$, and ν_i is primitive $3(q-1)$th root of unity, $i = 1, 2$.

As $\gcd(3(q-1),l)=1$, we can find a bijection θ from the set D to itself such that $\theta(\nu)=\nu^l$, for any $\nu\in D$, where D consists of all the primitive $3(q-1)th$ roots of unity of $F_{q^3}^*$. Therefore, there exists a unique element $\omega_i\in D$ such that $\nu_i^{-1}=\theta(\omega_i)=\omega_i^l$, i.e., $\omega_i^l\nu_i=1$, $i=1,2$.

From the above discussion, we have the following two lemmas.

Lemma 4.4. The irreducible factorization of $x^{3l}-\xi$ over F_q is given as follows:
(I) If $\gcd(f,3)=1$,

$$x^{3l}-\xi=\prod_{i=0}^{e}R_i(x),$$

where $R_i(x)=\widehat{M}_i(\omega_1x)\widehat{M}_i(\alpha\omega_1x)\widehat{M}_i(\alpha^2\omega_1x)$ for any $i=0,1,2,...,e$, with ω_1 is a primitive $3(q-1)$th root of unity and α is a primitive 3th root of unity.
(II) If $\gcd(f,3)=3$,

$$x^{3l}-\xi=P(x)\prod_{i=1}^{e}Q_i(x)U_i(x)Z_i(x),$$

where $P(x)=(x-\omega_1^{-1})(x-\alpha\omega_1^{-1})(x-\alpha^2\omega_1^{-1})$, $Q_i(x)=\widehat{A}_i(\omega_1x)\widehat{A}_{iq}(\alpha\omega_1x)\widehat{A}_{iq^2}(\alpha^2\omega_1x)$, $U_i(x)=\widehat{A}_i(\alpha\omega_1x)\widehat{A}_{iq}(\alpha^2\omega_1x)\widehat{A}_{iq^2}(\omega_1x)$, and $Z_i(x)=\widehat{A}_i(\alpha^2\omega_1x)\widehat{A}_{iq}(\omega_1x)\widehat{A}_{iq^2}(\alpha\omega_1x)$ for any $i=1,2,...,e$, with ω_1 is a primitive $3(q-1)$th root of unity, α is a primitive 3th root of unity.

Proof. (I) Ifgcd$(f,3)=1$, we get that C_0 and C_k,$1\le k\le e=\frac{\phi(l)}{f}$ are all the q^3-cyclotomic cosets modulo l. Let ν_1 be a root of $x^3-\xi$, and α be a primitive 3th root of unity. Then we have $\nu_1,\alpha\nu_1$, and $\alpha^2\nu_1$ are all the roots of $x^3-\xi$ over F_{q^3}, i.e.,

$$x^{3l}-\xi=(x^l-\nu_1)(x^l-\alpha\nu_1)(x^l-\alpha^2\nu_1).$$

From the above discussion, we know that there exists ω_1 such that $\omega_1^l\nu_1=1$, where ω_1 is a primitive $3(q-1)$th root of unity. Hence, we have $\omega_1^q=\alpha\omega_1$ or $\alpha^2\omega_1$. Next, we just consider $\omega_1^q=\alpha\omega_1$. However, for the case $\omega_1^q=\alpha^2\omega_1$, we see that the result is still right in the same way. As $\gcd(3,l)=1$, we know $l\equiv1(\mathrm{mod}\ 3)$ or $l\equiv2(\mathrm{mod}\ 3)$. When $l\equiv2(\mathrm{mod}\ 3)$, we have $(\alpha\omega_1)^l\alpha\nu_1=\alpha^{l+1}=1$ and $(\alpha^2\omega_1)^l\alpha^2\nu_1=\alpha^{2(l+1)}=1$. When $l\equiv1(\mathrm{mod}\ 3)$, we have $(\alpha^2\omega_1)^l\alpha\nu_1=\alpha^{2l+1}=1$ and $(\alpha\omega_1)^l\alpha^2\nu_1=\alpha^{l+2}=1$. Hence, there always exist $\omega_1,\alpha\omega_1$ and $\alpha^2\omega_1$ such that $\omega_1^l\nu_1=1$, $(\alpha\omega_1)^l\alpha\nu_1=1$ and $(\alpha^2\omega_1)^l\alpha^2\nu_1=1$ or $\omega_1^l\nu_1=1$, $(\alpha^2\omega_1)^l\alpha\nu_1=1$ and $(\alpha\omega_1)^l\alpha^2\nu_1=1$. Further, by Lemma 4.2, we get

$$x^{3l}-\xi=\prod_{i=0}^{e}\widehat{M}_i(\omega_1x)\widehat{M}_i(\alpha\omega_1x)\widehat{M}_i(\alpha^2\omega_1x),$$

which is the monic irreducible factorization of $x^{3l}-\xi$ over F_{q^3}. And we have $\widehat{M}_i(\omega_1x)=\prod_{k\in C_i}(x-\omega_1^{-1}\eta^k)$, $\widehat{M}_i(\alpha\omega_1x)=\prod_{k\in C_i}(x-\alpha^2\omega_1^{-1}\eta^k)$ and $\widehat{M}_i(\alpha^2\omega_1x)=\prod_{k\in C_i}(x-\alpha\omega_1^{-1}\eta^k)$. Obviously, when k runs over C_i, $\omega_1^{-1}\eta^k$ gives all the roots of $\widehat{M}_i(\omega_1x)$. As $\omega_1^q=\alpha\omega_1$ and $kq,kq^2\in C_i$, we have $(\omega_1^{-1}\eta^k)^q=\alpha^2\omega_1^{-1}\eta^{kq}$ and $(\omega_1^{-1}\eta^k)^{q^2}=\alpha\omega_1^{-1}\eta^{kq^2}$, which gives a root of $\widehat{M}_i(\alpha\omega_1x)$ and $\widehat{M}_i(\alpha^2\omega_1x)$ respectively. Therefore, it's easy to deduce that $\widehat{M}_i(\omega_1x)\widehat{M}_i(\alpha^2\omega_1x)$ $\widehat{M}_i(\alpha^2\omega_1x)$ is a irreducible polynomial over F_q.
(II) When $\gcd(f,3)=3$, we have that A_0,A_k,A_{kq},A_{kq^2} consist of all the distinct q^3-cyclotomic cosets modulo l, where $1\le k\le e$. Then, the irreducible factorization of x^l-1 over F_{q^3} is given by

$$x^l-1=A_0(x)A_1(x)A_q(x)A_{q^2}(x)A_2(x)A_{2q}(x)A_{2q^2}(x)...A_e(x)A_{eq}(x)A_{eq^2}(x).$$

Next, in the same way as (I), we can proved the conclusion (II) holds. □

Using arguments similar to the proof in Lemma 4.3, we have the following lemma, and we omit its proof here.

Lemma 4.5. The irreducible factorization of $x^{3l}-\xi^2$ over F_q is given as follows:
(I) If $\gcd(f,3)=1$,

$$x^{3l}-\xi^2=\prod_{i=0}^{e}R_i^{'}(x),$$

where $R_i^{'}(x)=\widehat{M}_i(\omega_2 x)\widehat{M}_i(\alpha\omega_2 x)\widehat{M}_i(\alpha^2\omega_2 x)$ for any $i=0,1,2,...,e$, with ω_2 is a primitive $3(q-1)$th root of unity and α is a primitive 3th root of unity.
(II) If $\gcd(f,3)=3$,

$$x^{3l}-\xi^2=P^{'}(x)\prod_{i=1}^{e}Q_i^{'}(x)U_i^{'}(x)Z_i^{'}(x),$$

where $P^{'}(x)=(x-\omega_2^{-1})(x-\alpha\omega_2^{-1})(x-\alpha^2\omega_2^{-1})$, $Q_i^{'}(x)=\widehat{A}_i(\omega_2 x)\widehat{A}_{iq}(\alpha\omega_2 x)\widehat{A}_{iq^2}(\alpha^2\omega_2 x)$, $U_i^{'}(x)=\widehat{A}_i(\alpha\omega_2 x)\widehat{A}_{iq}(\alpha^2\omega_2 x)\widehat{A}_{iq^2}(\omega_2 x)$, and $Z_i^{'}(x)=\widehat{A}_i(\alpha^2\omega_2 x)\widehat{A}_{iq}(\omega_2 x)\widehat{A}_{iq^2}(\alpha\omega_2 x)$ for any $i=1,2,...,e$, with ω_2 is a primitive $3(q-1)$th root of unity and α is a primitive 3th root of unity.

Combining Lemmas 4.3, 4.4 and 4.5, we easily obtain the next result.

Theorem 4.6. Let $\gcd(q-1,3lp^s)=3$ and $\alpha=\xi^{\frac{q-1}{3}}$. For any element λ of F_q^* and $\lambda-$constacyclic code C of length $3lp^s$ over F_q, one of the following cases holds:
(I) If $\lambda\in\langle\xi^3\rangle$, then there exists some element $c\in F_q^*$ such that $c^{3lp^s}\lambda=1$, and we have

$$C=\langle\prod_{i=0}^{e}\widehat{M}_i(cx)^{\varepsilon_i}\widehat{M}_i(cb_{\frac{q-1}{3}}x)^{\sigma_i}\widehat{M}_i(cb_{\frac{2(q-1)}{3}}x)^{\tau_i}\rangle,$$

$$C^{\perp}=\langle\prod_{i=0}^{e}\widehat{M}_{-i}(c^{-1}x)^{p^s-\varepsilon_i}\widehat{M}_{-i}(c^{-1}b_{\frac{q-1}{3}}^{-1}x)^{p^s-\sigma_i}\widehat{M}_{-i}(c^{-1}b_{\frac{2(q-1)}{3}}^{-1}x)^{p^s-\tau_i}\rangle,$$

where $0\le\varepsilon_i,\sigma_i,\tau_i\le p^s$ for any $i=0,1,2,...,e$.
(II) If $\lambda\in\xi^{p^s}\langle\xi^3\rangle$, then there exists some element $c_1\in F_q^*$ such that $c_1^{3lp^s}\lambda=\xi^{p^s}$, and one of the following holds:
(i) If $\gcd(f,3)=1$,

$$C=\langle\prod_{i=0}^{e}\widehat{R}_i(c_1x)^{\varepsilon}\rangle,$$

$$C^{\perp}=\langle\prod_{i=0}^{e}\widehat{R}_{-i}(c_1^{-1}x)^{p^s-\varepsilon}\rangle,$$

where $0\le\varepsilon\le p^s$, $R_{-i}(x)=\widehat{M}_{-i}(\omega_1^{-1}x)\widehat{M}_{-i}(\alpha^2\omega_1^{-1}x)\widehat{M}_{-i}(\alpha\omega_1^{-1}x)$ for any $i=0,1,2,...,e$.
(ii) If $\gcd(f,3)=3$,

$$C=\langle\widehat{P}(c_1x)^{\varepsilon}\prod_{i=1}^{e}\widehat{Q}_i(c_1x)^{\varepsilon_i}\widehat{U}_i(c_1x)^{\sigma_i}\widehat{Z}_i(c_1x)^{\tau_i}\rangle,$$

$$C^{\perp} = \langle \widehat{P}^*(c_1^{-1}x)^{p^s-\varepsilon} \prod_{i=1}^{e} \widehat{Q}_{-i}(c_1^{-1}x)^{p^s-\varepsilon_i} \widehat{U}_{-i}(c_1^{-1}x)^{p^s-\sigma_i} \widehat{Z}_{-i}(c_1^{-1}x)^{p^s-\tau_i} \rangle,$$

where $0 \leq \varepsilon_i, \sigma_i, \tau_i \leq p^s$, $P^*(x) = (x-\omega_1)(x-\alpha^2\omega_1)(x-\alpha\omega_1)$, $Q_{-i}(x) = \widehat{A}_{-i}(\omega_1^{-1}x) \cdot \widehat{A}_{-iq}(\alpha^2\omega_1^{-1}x)\widehat{A}_{-iq^2}(\alpha\omega_1^{-1}x)$, $U_{-i}(x) = \widehat{A}_{-i}(\alpha^2\omega_1^{-1}x)\widehat{A}_{-iq}(\alpha\omega_1^{-1}x)\widehat{A}_{-iq^2}(\omega_1^{-1}x)$, and $Z_{-i}(x) = \widehat{A}_{-i}(\alpha\omega_1^{-1}x)\widehat{A}_{-iq}(\omega_1^{-1}x)\widehat{A}_{-iq^2}(\alpha^2\omega_1^{-1}x)$, for any $i = 1, 2, ..., e$.

(III) If $\lambda \in \xi^{2p^s}\langle \xi^3 \rangle$, then there exists some element $c_2 \in F_q^*$ such that $c_2^{3lp^s}\lambda = \xi^{2p^s}$, and one of the following holds:

(i) If $\gcd(f, 3) = 1$,

$$C = \langle \prod_{i=0}^{e} \widehat{R'}_i(c_2x)^{\varepsilon_i} \rangle,$$

$$C^{\perp} = \langle \prod_{i=0}^{e} \widehat{R'}_{-i}(c_2^{-1}x)^{p^s-\varepsilon_i} \rangle,$$

where $0 \leq \varepsilon \leq p^s$, $R'_{-i}(x) = \widehat{M}_{-i}(\omega_2^{-1}x)\widehat{M}_{-i}(\alpha^2\omega_2^{-1}x)\widehat{M}_{-i}(\alpha\omega_2^{-1}x)$ for any $i = 0, 1, 2, ..., e$.

(ii) If $\gcd(f, 3) = 3$,

$$C = \langle \widehat{P'}(c_2x)^{\varepsilon} \prod_{i=1}^{e} \widehat{Q'}_i(c_2x)^{\varepsilon_i} \widehat{U'}_i(c_2x)^{\sigma_i} \widehat{Z'}_i(c_2x)^{\tau_i}) \rangle,$$

$$C^{\perp} = \langle \widehat{P'}^*(c_2^{-1}x)^{p^s-\varepsilon} \prod_{i=1}^{e} \widehat{Q'}_{-i}(c_2^{-1}x)^{p^s-\varepsilon_i} \widehat{U'}_{-i}(c_2^{-1}x)^{p^s-\sigma_i} \widehat{Z'}_{-i}(c_2^{-1}x)^{p^s-\tau_i}) \rangle,$$

where $0 \leq \varepsilon_i, \sigma_i, \tau_i \leq p^s$, $P'^*(x) = (x-\omega_2)(x-\alpha^2\omega_2)(x-\alpha\omega_2)$, $Q'_{-i}(x) = \widehat{A}_{-i}(\omega_2^{-1}x) \cdot \widehat{A}_{-iq}(\alpha^2\omega_2^{-1}x)\widehat{A}_{-iq^2}(\alpha\omega_2^{-1}x)$, $U'_{-i}(x) = \widehat{A}_{-i}(\alpha^2\omega_2^{-1}x)\widehat{A}_{-iq}(\alpha\omega_2^{-1}x)\widehat{A}_{-iq^2}(\omega_2^{-1}x)$, and $Z'_{-i}(x) = \widehat{A}_{-i}(\alpha\omega_2^{-1}x)\widehat{A}_{-iq}(\omega_2^{-1}x)\widehat{A}_{-iq^2}(\alpha^2\omega_2^{-1}x)$ for any $i = 1, 2, ..., e$.

4.3 All constacyclic codes of length $3lp^s$ over F_q when $d = l$

Let $d = \gcd(q-1, 3lp^s) = l$, i.e., $l|(q-1)$ and $\gcd(q-1, 3) = 1$. Then $\eta = \xi^{\frac{q-1}{l}} \in F_q^*$ is a primitive lth root of unity. Therefore, we have the following lemma.

Lemma 4.7. Assume that $\gcd(q-1, 3lp^s) = l$, and let $\eta = \xi^{\frac{q-1}{l}}$. Then, the irreducible factorization of $x^{3lp^s} - 1$ over F_q is given by:

$$x^{3lp^s} - 1 = \prod_{k=0}^{l-1} (x-\eta^k)^{p^s}(x^2 + \eta^k x + \eta^{2k})^{p^s}.$$

Proof. As $\gcd(q-1, 3) = 1$, there exists no primitive 3th root of unity in F_q, which implies $x^2 + x + 1$ is irreducible. By this, we can deduce that $x^2 + \eta^k x + \eta^{2k}$ is irreducible, for any $k = 1, 2, ..., l$. Otherwise $x^2 + \eta^k x + \eta^{2k}$ is reducible, then we have $\eta^{-2k}(x^2 + \eta^k x + \eta^{2k}) = (\eta^{-k}x)^2 + \eta^{-k}x + 1$ is reducible. By substituting x for $\eta^{-k}x$ in the above polynomial, we have that $x^2 + x + 1$ is reducible, which is a contradiction. Since $\eta = \xi^{\frac{q-1}{l}}$ be a primitive lth root

of unity, then η^3 is also a primitive lth root of unity. Hence, the irreducible factorization of $x^{3lp^s}-1$ over F_q is given by

$$x^{3lp^s}-1=(x^{3l}-1)^{p^s}=\prod_{k=0}^{l-1}(x^3-\eta^{3k})^{p^s}=\prod_{k=0}^{l-1}(x-\eta^k)^{p^s}(x^2+\eta^k x+\eta^{2k})^{p^s}.$$

□

Lemma 4.8. Assume that $\gcd(q-1,3lp^s)=l$, then the irreducible factorization of $x^{3lp^s}-\xi^{jp^s}$, $1\leq j\leq l-1$, over F_q is given by:
(I) When $\gcd(3,j)=3$, let $j=3k$, for some integer k. Then we have

$$x^{3lp^s}-\xi^{jp^s}=(x^l-\xi^k)^{p^s}(x^{2l}+\xi^k x^l+\xi^{2k})^{p^s}.$$

(II) When $\gcd(3,j)=1$, there must exist some integer i, $1\leq i\leq q-1$, such that $3i=j+q-1$ or $3i=j+2(q-1)$. Then we have

$$x^{3lp^s}-\xi^{jp^s}=(x^l-\xi^i)^{p^s}(x^{2l}+\xi^i x^l+\xi^{2i})^{p^s}.$$

Proof. (I) Obviously, $\gcd(l,k)=1$. From Lemma 2.3, it is very easy to verify that $x^l-\xi^k$ is irreducible. By the proof of Lemma 4.5, we see that $x^2+\xi^k x+\xi^{2k}$ is irreducible over F_q. Now, we suppose that δ is any root of $x^2+\xi^k x+\xi^{2k}$ in some extended field of F_q, and e is the order of δ. Then, we have $\delta^3=\xi^{3k}$. Further, we deduce that $\frac{e}{(e,3)}=\frac{q-1}{(q-1,3k)}$, i.e., $\frac{e}{(e,3)}=\frac{q-1}{(q-1,k)}$, as $\gcd(q-1,3)=1$. By the reduction again, we get $e=\frac{(q-1)(e,3)}{(q-1,k)}$. From Lemma 2.4, we can verify that $x^{2l}+\xi^k x^l+\xi^{2k}$ is irreducible.

(II) When $(3,j)=1$, we get that $x^3-\xi^j$, $1\leq j\leq l-1$, are all reducible, from Lemma 2.3. Therefore, there must exist some $\xi^i\in F_q^*$, which is a root of $x^3-\xi^j$, for any $j=1,2,...l-1$. Then, $\xi^{3i}-\xi^j=0$, i.e., $\xi^{3i}=\xi^j$. As $1\leq i\leq q-1$ and $1\leq j\leq l-1$, we deduce that $3i=j+q-1$ or $3i=j+2(q-1)$ and $\gcd(l,i)=1$. Next, working similar to the proof of (I), we get that the results (i) and (ii) hold. □

According to Lemma 4.7 and Lemma 4.8, we get the following theorem immediately.

Theorem 4.9. Assume that $\gcd(q-1,3lp^s)=l$, and let $\eta=\xi^{\frac{q-1}{l}}$. For any element λ of F_q^* and $\lambda-$constacyclic code C of length $3lp^s$ over F_q, one of the following cases holds:
(I) If $\lambda\in\langle\xi^l\rangle$, then there exists $c_1\in F_q^*$ such that $c_1^{3lp^s}\lambda=1$, and we have

$$C=\langle\prod_{k=0}^{l-1}(x-c_1^{-1}\eta^k)^{\varepsilon_k}(x^2+c_1^{-1}\eta^k x+c_1^{-2}\eta^{2k})^{\tau_k}\rangle,$$

$$C^{\perp}=\langle\prod_{k=0}^{l-1}(x-c_1\eta^{-k})^{p^s-\varepsilon_k}(x^2+c_1\eta^{-k}x+c_1^2\eta^{-2k})^{p^s-\tau_k}\rangle,$$

where $0\leq\varepsilon_k,\tau_k\leq p^s$, for any $k=0,1,2,...,l-1$.
(II) If $\lambda\in\xi^{jp^s}\langle\xi^l\rangle$, $1\leq j\leq l-1$, then there exists $c_2\in F_q^*$ such that $c_2^{3lp^s}\lambda=\xi^{jp^s}$, and one of the following holds:
(i) When $(3,j)=3$, let $j=3k$, for some integer k. Then we have

$$C=\langle(x^l-c_2^{-l}\xi^k)^{\varepsilon_k}(x^{2l}+c_2^{-l}\xi^k x^l+c_2^{-2l}\xi^{2k})^{\tau_k}\rangle,$$

$$C^{\perp} = \langle (x^l - c_2^l \xi^{-k})^{p^s - \varepsilon_k} (x^{2l} + c_2^l \xi^{-k} x^l + c_2^{2l} \xi^{-2k})^{p^s - \tau_k} \rangle,$$

where $0 \leq \varepsilon_k, \tau_k \leq p^s$.

(ii) When $\gcd(3, j) = 1$, there must exist some integer i, $1 \leq i \leq q-1$, such that $3i = j+q-1$ or $3i = j + 2(q-1)$. Then we have

$$C = \langle (x^l - c_2^{-l} \xi^i)^{\varepsilon_i} (x^{2l} + c_2^{-l} \xi^i x^l + c_2^{-2l} \xi^{2i})^{\tau_i} \rangle,$$

$$C^{\perp} = \langle (x^l - c_2^l \xi^{-i})^{p^s - \varepsilon_i} (x^{2l} + c_2^l \xi^{-i} x^l + c_2^{2l} \xi^{-2i})^{p^s - \tau_i} \rangle.$$

where $0 \leq \varepsilon_i, \tau_i \leq p^s$.

4.4 All constacyclic codes of length $3lp^s$ over F_q when $d = 3l$

In this subsection, we assume that $d = \gcd(3lp^s, q-1) = 3l$, namely $3l|(q-1)$. Clearly, there exists an element $\gamma = \xi^{\frac{q-1}{3l}} \in F_q^*$, which is a primitive $3l$th root of unity. Further, due to $l|(q-1)$, and $3|(q-1)$, it is easy to know that $\eta = \xi^{\frac{q-1}{l}}$ and $\beta = \xi^{\frac{q-1}{3}}$ are primitive lth and 3th roots of unity respectively.

From Lemma 3.1, we get that the $F_q^* = \langle \xi \rangle = \langle \xi^{3l} \rangle \cup \xi^{p^s} \langle \xi^{3l} \rangle \cup \xi^{2p^s} \langle \xi^{3l} \rangle \cup ... \cup \xi^{(3l-1)p^s} \langle \xi^{3l} \rangle$. Therefore, any element λ of F_q^* belongs to exactly one of the cosets, i.e., there is a unique integer j, $0 \leq j \leq 3l-1$, such that $\lambda \in \xi^{jp^s} \langle \xi^{3l} \rangle$, namely $\lambda-$constacyclic codes are equivalent to $\xi^{jp^s}-$constacyclic codes. Hence, we just need to determine $\xi^{jp^s}-$constacyclic codes, where $0 \leq j \leq 3l-1$.

Lemma 4.10. Let $d = \gcd(3lp^s, q-1) = 3l$ and $\gamma = \xi^{\frac{q-1}{3l}}$. Then irreducible factorization of $x^{3lp^s} - 1$ over F_q is given as follows:

$$x^{3lp^s} - 1 = \prod_{i=0}^{3l-1} (x - \gamma^i)^{p^s}.$$

Lemma 4.11. Let $\eta = \xi^{\frac{q-1}{l}}$ and $\beta = \xi^{\frac{q-1}{3}}$. Then the irreducible factorization of $x^{3lp^s} - \xi^{jp^s}$ over F_q is given as follows:
(I) When $\gcd(3l, j) = l$, we have

$$x^{3lp^s} - \xi^{jp^s} = (x^{3l} - \xi^{tl})^{p^s} = \prod_{i=0}^{l-1} (x^3 - \xi^t \eta^i)^{p^s},$$

where $t = 1$ or 2.
(II) When $\gcd(3l, j) = 3$, we have

$$x^{3lp^s} - \xi^{jp^s} = (x^{3l} - \xi^{3k})^{p^s} = \prod_{i=0}^{2} (x^l - \xi^k \beta^i)^{p^s},$$

where k is some integer such that $j = 3k$.
(III) Otherwise, we can see that $\gcd(3l, j) = 1$. Then we have

$$x^{3lp^s} - \xi^{jp^s} = (x^{3l} - \xi^j)^{p^s}.$$

Proof. (I) As $\gcd(3l, j) = l$ and $1 \leq j \leq 3l-1$, we have $j = tl$, where $t = 1, 2$. Obviously, $\eta = \xi^{\frac{q-1}{l}}$ is a primitive $l-$th root of unity in F_q. Therefore, we get

$$x^{3lp^s} - \xi^{jp^s} = (x^{3l} - \xi^{tl})^{p^s} = \prod_{i=0}^{l-1}(x^3 - \xi^t\eta^i)^{p^s},$$

Next, we prove that the polynomial $x^3 - \xi^t\eta^i$, for any $i = 0, 1, 2, ..., l-1$, is irreducible in $F_q[x]$. Firstly, we know that the multiplicative order of $\xi^t\eta^i = \xi^{t+\frac{i(q-1)}{l}}$, $t = 1, 2$, is $e_i = \frac{q-1}{(q-1, t+\frac{i(q-1)}{l})}$. As $3|(q-1)$, $\gcd(3, t) = 1$ and $\gcd(3, l) = 1$, we get that $(3, t + \frac{i(q-1)}{l}) = 1$. Thus, 3 divides e_i but not $\frac{q-1}{e_i} = (q-1, t + \frac{i(q-1)}{l})$. From Lemma 2.3, we get the polynomial $x^3 - \xi^t\eta^i$, for any $i = 0, 1, 2, ..., l-1$, is irreducible in $F_q[x]$. In the same way, we have (II) and (III) hold. □

In the following theorem, we determine all constacyclic codes of length $3lp^s$ over F_q and their dual codes, when $d = \gcd(3lp^s, q-1) = 3l$.

Theorem 4.12. Assume that $\gcd(3lp^s, q-1) = 3l$, let $\gamma = \xi^{\frac{q-1}{3l}}, \eta = \xi^{\frac{q-1}{l}}$ and $\beta = \xi^{\frac{q-1}{3}}$ be primitive $3l$th, lth and 3th root of unity in F_q respectively. For any element λ of F_q^* and $\lambda-$ constcyclic codes C of length $3lp^s$ over F_q. One of the following holds:
(I) If $\lambda \in \langle \xi^{3l} \rangle$, then there exists $d_1 \in F_q^*$ such that $d_1^{3lp^s}\lambda = 1$, and we have

$$C = \langle \prod_{i=0}^{3l-1}(x - d_1^{-1}\gamma^i)^{\varepsilon_i} \rangle,$$

$$C^{\perp} = \langle \prod_{i=0}^{3l-1}(x - d_1\gamma^{-i})^{p^s-\varepsilon_i} \rangle,$$

where $0 \leq \varepsilon_i \leq p^s$, for any $i = 0, 1, 2, ..., 3l-1$.
(II) If $\lambda \in \xi^{jp^s}\langle \xi^{3l} \rangle$, $1 \leq j \leq 3l-1$, then there exists $d_2 \in F_q^*$ such that $d_2^{3lp^s}\lambda = \xi^{jp^s}$, and one of the following holds:
(i) When $\gcd(3l, j) = l$, we have

$$C = \langle \prod_{i=0}^{l-1}(x^3 - d_2^{-3}\xi^t\eta^i)^{\varepsilon_i} \rangle,$$

$$C^{\perp} = \langle \prod_{i=0}^{l-1}(x^3 - d_2^3\xi^{-t}\eta^{-i})^{p^s-\varepsilon_i} \rangle,$$

where $0 \leq \varepsilon_i \leq p^s$, for any $i = 0, 1, 2, ..., l-1$.
(ii) When $\gcd(3l, j) = 3$, we have

$$C = \langle \prod_{i=0}^{2}(x^l - d_2^{-l}\xi^k\beta^i)^{\varepsilon_i} \rangle,$$

$$C^{\perp} = \langle \prod_{i=0}^{2}(x^l - d_2^l\xi^{-k}\beta^{-i})^{p^s-\varepsilon_i} \rangle,$$

where $0 \leq \varepsilon_i \leq p^s$ for $i = 0, 1, 2$, and $j = 3k$.

(iii) When $\gcd(3l, j) = 1$, we have

$$C = \langle (x^{3l} - d_2^{-3l}\xi^j)^{\varepsilon} \rangle,$$

$$C^{\perp} = \langle (x^{3l} - d_2^{3l}\xi^{-j})^{p^s - \varepsilon} \rangle,$$

where $0 \leq \varepsilon \leq p^s$.

5 All self-dual cyclic codes of length $3lp^s$ over F_q

In Section 4, we gave the generator polynomials of all the constacyclic codes and their dual codes of length $3lp^s$ over F_q. We will determine all the self-dual cyclic codes of length $3lp^s$ over F_q in detail, in this section.

It is well known that there exist self-dual cyclic codes of length N over F_q if and only if N is even and the characteristic of F_q is $p = 2$ [10, 11]. Therefore, we get that self-dual cyclic codes of length $3lp^s$ over F_q exist only when $p = 2$.

Let $x^{3lp^s} - 1 = (x^{3l} - 1)^{p^s} = f_1(x)^{p^s} f_2(x)^{p^s} \cdots f_a(x)^{p^s} h_1(x)^{p^s} h_1^*(x)^{p^s} \cdots h_b(x)^{p^s} h_b^*(x)^{p^s}$ be the irreducible factorization of $x^{3lp^s} - 1$, where $f_1(x), f_2(x), ..., f_a(x)$ are monic irreducible self-reciprocal polynomials over F_q, $h_j(x)$ and its reciprocal polynomial $h_j^*(x)$, $1 \leq j \leq b$, are also monic irreducible polynomials over F_q. Hence, for any cyclic code $C = \langle g(x) \rangle$ of length $3lp^s$ over F_q, we suppose that

$$g(x) = f_1(x)^{\tau_1} f_2(x)^{\tau_2} \cdots f_a(x)^{\tau_a} h_1(x)^{\delta_1} h_1^*(x)^{\sigma_1} \cdots h_b(x)^{\delta_b} h_b^*(x)^{\sigma_b},$$

where $0 \leq \tau_i, \delta_j, \sigma_j \leq p^s$, for any $i = 1, 2, ..., a$, and $j = 1, 2, ..., b$. Then, we have

$$h(x) = f_1(x)^{p^s - \tau_1} f_2(x)^{p^s - \tau_2} \cdots f_a(x)^{p^s - \tau_a} h_1(x)^{p^s - \delta_1} h_1^*(x)^{p^s - \sigma_1} \cdots h_b(x)^{p^s - \delta_b} h_b^*(x)^{p^s - \sigma_b}.$$

Therefore,

$$h^*(x) = f_1(x)^{p^s - \tau_1} f_2(x)^{p^s - \tau_2} \cdots f_a(x)^{p^s - \tau_a} h_1(x)^{p^s - \sigma_1} h_1^*(x)^{p^s - \delta_1} \cdots h_b(x)^{p^s - \sigma_b} h_b^*(x)^{p^s - \delta_b}.$$

If C is a self-dual cyclic code, we get the following theorem.

Theorem 5.1. With the above notations, we have that C is a self-dual cyclic code if and only if $2\tau_i = p^s$, $1 \leq i \leq a$, and $\delta_j + \sigma_j = p^s$, $1 \leq j \leq b$.

Proof. C is a self-dual cyclic code if and only if $g(x) = h^*(x)$, i.e., $2\tau_i = p^s$, $1 \leq i \leq a$, and $\delta_j + \sigma_j = p^s$, $1 \leq j \leq b$. □

According to this theorem, we see that it is enough to determine the irreducible factorization of $x^{3lp^s} - 1$ as above. And if we do this, we can give all the self-dual cyclic codes immediately.

Similar to the definition of reciprocal polynomial, we give the following definition.

Definition 5.2. Let $C_s = \{s, sq, ..., sq^{f-1}\}$ be any $q-$cyclotomic coset modulo l, then

$$C_s^* = \{-s, -sq, ..., -sq^{f-1}\}$$

is said to be the reciprocal coset of C_s. The coset C_s is called self-reciprocal if $C_s = C_s^*$.

Obviously, C_s^* is still a $q-$cyclotomic coset modulo l. And the reciprocal polynomial of the minimal polynomial of C_s is the minimal polynomial of C_s^*. Hence, the minimal polynomial of C_s is also self-reciprocal if C_s is self-reciprocal.

Lemma 5.3. Assume that $q \equiv 1(\text{mod } 3)$. For the $q-$cyclotomic cosets, which have been described in Lemma 2.1, one of the following holds:
(I) If $f = \text{ord}_l(q)$ is even, we have

$$B_0^* = B_0, B_l^* = B_{-l}, B_{g^k}^* = B_{-g^k}, B_{3g^k}^* = B_{3g^k},$$

where $0 \leq k \leq e-1$.
(II) If $f = \text{ord}_l(q)$ is odd, we have

$$B_0^* = B_0, B_l^* = B_{-l}, B_{g^k}^* = B_{-g^k}, B_{3g^{k'}}^* = B_{-3g^{k'}}$$

where $\{B_{3g^k}\} = \{B_{3g^{k'}}\} \bigcup \{B_{-3g^{k'}}\}$ and $0 \leq k \leq e-1$, $0 \leq k^{'} \leq \frac{e}{2} - 1$.

Proof. (I) Obviously, we only need to prove $B_{3g^k}^* = B_{3g^k}$. If $f = \text{ord}_l(q)$ is even, we deduce that $q^{\frac{f}{2}} \equiv -1(\text{mod } l)$. According to this, we get there exist i, j, $0 \leq i, j \leq f-1$, and $|j-i| = \frac{f}{2}$, such that $3g^k q^i \equiv -3g^k q^j (\text{mod } 3l)$, for any $3g^k q^i \in B_{3g^k}$, $0 \leq k \leq e-1$. Therefore, we have $B_{3g^k}^* = B_{3g^k}$, for any $k = 0, 1, ..., e-1$.

(II) In the same way as Lemma 2.1, we get that $B_0, B_l, B_{-l}, B_{g^k}, B_{-g^k}, B_{3g^{k'}}$ and $B_{-3g^{k'}}$, $0 \leq k \leq e-1$, $0 \leq k^{'} \leq \frac{e}{2} - 1$, are all the distinct $q-$cyclotomic cosets modulo $3l$. The next result is obvious. □

Lemma 5.4. Let $q \equiv 2(\text{mod } 3)$ and f be even. For the $q-$cyclotomic cosets, which have been described in Lemma 2.1, one of the following holds:
(I) When $f = 2t$ and t is even, we have

$$B_0^* = B_0, B_l^* = B_l, B_{g^k}^* = B_{-g^k}, B_{3g^k}^* = B_{3g^k},$$

where $\{B_{g^{k'}}\} = \{B_{g^k}\} \bigcup \{B_{-g^k}\}$,$0 \leq k \leq e-1$ and $0 \leq k^{'} \leq 2e-1$.
(II) When $f = 2t$ and t is odd, we have

$$B_0^* = B_0, B_l^* = B_l, B_{g^{k'}}^* = B_{g^{k'}}, B_{3g^k}^* = B_{3g^k},$$

where $0 \leq k \leq e-1$ and $0 \leq k^{'} \leq 2e-1$.

Proof. (I) In the same way as Lemma 2.1, we get $B_0, B_l, B_{g^k}, B_{-g^k}$ and B_{3g^k} , $0 \leq k \leq e-1$, are all the distinct $q-$cyclotomic cosets modulo $3l$. Next, We first prove that $B_l^* = B_l$, i.e., $\{l, lq\}^* = \{l, lq\}$. As $q \equiv 2(\text{mod } 3)$, i.e., $q \equiv -1(\text{mod } 3)$, then $lq \equiv -l(\text{mod } 3)$. Since $\gcd(3, l) = 1$, we have $lq \equiv -l(\text{mod } 3l)$, which implies $B_l^* = B_l$. Otherwise, by the conclusion (I) of Lemma 5.3, we get the other results immediately.

(II) According to (I), it is obvious that we only need to prove $B_{g^{k'}}^* = B_{g^{k'}}$. As t is odd, we have $q^t \equiv -1(\text{mod } 3)$. Since $q^t \equiv -1(\text{mod } l)$ and $\gcd(3, l) = 1$, we get $q^t \equiv -1(\text{mod } 3l)$. Then, we deduce that there exist i, j, $0 \leq i, j \leq f-1$, and $|j-i| = t$, such that $g^{k^{'}} q^i \equiv -g^{k^{'}} q^j (\text{mod } 3l)$, for any $g^{k^{'}} q^i \in B_{k'}$, $0 \leq k^{'} \leq 2e-1$, i.e., $B_{g^{k'}}^* = B_{g^{k'}}$. □

Lemma 5.5. Let $q \equiv 2(\text{mod } 3)$ and f be odd. For the $q-$cyclotomic cosets, which have been described in Lemma 2.1, we have

$$B_0^* = B_0, B_l^* = B_l, B_{g^{k'}}^* = B_{-g^{k'}}, B_{3g^{k'}}^* = B_{-3g^{k'}},$$

where $\{B_{g^k}\} = \{B_{g^{k'}}\} \bigcup \{B_{-g^{k'}}\}$, $\{B_{3g^k}\} = \{B_{3g^{k'}}\} \bigcup \{B_{-3g^{k'}}\}$ and $0 \leq k \leq e-1$, $0 \leq k^{'} \leq \frac{e}{2} - 1$.

From the above lemmas, we can give all the self-dual cyclic codes of length $3 \cdot 2^s l$ over F_{2^m} and their enumeration in the following theorem.

Theorem 5.6. Let $l \neq 3$ be an odd prime, $p = 2$, $f = \text{ord}_l(2^m)$, and $e = \frac{l-1}{f}$. Then, for cyclic self-dual codes of length $3 \cdot 2^s l$ over F_{2^m}, we have
(I) When $q \equiv 1(\text{mod } 3)$, one of the following hold:
(i) If $f = \text{ord}_l(q)$ is even, then there exist $(2^s+1)^{e+1}$ cyclic self-dual codes of length $3 \cdot 2^s l$ over F_{2^m} given by

$$\langle (x-1)^{2^{s-1}} B_l(x)^{\delta} B_{-l}(x)^{2^s-\delta} \prod_{k=0}^{e-1} B_{g^k}(x)^{\delta_k} B_{-g^k}(x)^{2^s-\delta_k} B_{3g^k}(x)^{2^{s-1}} \rangle,$$

where $0 \leq \delta, \delta_k \leq 2^s$, for any $0 \leq k \leq e-1$.
(ii) If $f = \text{ord}_l(q)$ is odd, then there exist $(2^s+1)^{\frac{3e}{2}+1}$ cyclic self-dual codes of length $3 \cdot 2^s l$ over F_{2^m} given by

$$\langle (x-1)^{2^{s-1}} B_l(x)^{\delta} B_{-l}(x)^{2^s-\delta} \prod_{k=0}^{e-1} B_{g^k}(x)^{\delta_k} B_{-g^k}(x)^{2^s-\delta_k} \prod_{k'=0}^{\frac{e}{2}-1} B_{3g^{k'}}(x)^{\sigma_{k'}} B_{-3g^{k'}}(x)^{2^s-\sigma_{k'}} \rangle,$$

where $0 \leq \delta, \delta_k, \sigma_{k'} \leq 2^s$, for any $0 \leq k \leq e$ and $0 \leq k^{'} \leq \frac{e}{2} - 1$.
(II) When $q \equiv 2(\text{mod } 3)$, we have
(i) If $f = 2t$ and t is even, then there exist $(2^s+1)^e$ cyclic self-dual codes of length $3 \cdot 2^s l$ over F_{2^m} given by

$$\langle (x-1)^{2^{s-1}} B_l(x)^{2^{s-1}} \prod_{k=0}^{e-1} B_{g^k}(x)^{\delta_k} B_{-g^k}(x)^{2^s-\delta_k} B_{3g^k}(x)^{2^{s-1}} \rangle,$$

where $0 \leq \delta_k \leq 2^s$, for any $0 \leq k \leq e-1$.
(ii) If $f = 2t$ and t is odd, then there exists only one cyclic self-dual code of length $3 \cdot 2^s l$ over F_{2^m} given by

$$\langle (x-1)^{2^{s-1}} B_l(x)^{2^{s-1}} \prod_{k'=0}^{2e-1} B_{g^{k'}}(x)^{2^{s-1}} \prod_{k=0}^{e-1} B_{3g^k}(x)^{2^{s-1}} \rangle.$$

(iii) If f is odd, then there exist $(2^s+1)^e$ cyclic self-dual codes of length $3 \cdot 2^s l$ over F_{2^m} given by

$$\langle (x-1)^{2^{s-1}} B_l(x)^{2^{s-1}} \prod_{k'=0}^{\frac{e}{2}-1} B_{g^{k'}}(x)^{\delta_{k'}} B_{-g^{k'}}(x)^{2^s-\delta_{k'}} B_{3g^{k'}}(x)^{\sigma_{k'}} B_{-3g^{k'}}(x)^{2^s-\sigma_{k'}} \rangle.$$

where $0 \leq \delta_{k'}, \sigma_{k'} \leq 2^s$, for any $0 \leq k^{'} \leq \frac{e}{2} - 1$.

References

[1] G. Castagnoli, J.L. Massey, P.A. Schoeller, N. von Seemann, On repeated-root cyclic codes, IEEE Trans. Inf. Theory 37 (1991) 337-342.

[2] J.H. van Lint, Repeated-root cyclic codes, IEEE Trans. Inf. Theory 37 (1991) 343-345.

[3] H.Q. Dinh, Repeated-root constacyclic codes of length $2p^s$, Finite Fields Appl. 18 (2012) 940-950.

[4] H.Q. Dinh, Structure of repeated-root constacyclic codes of length $3p^s$ and their duals, Discrete Math. 313 (2013) 983-991.

[5] H.Q. Dinh, Structure of repeated-root cyclic codes and negacyclic codes of length $6p^s$ and their duals, Contemp. Math. 609 (2014) 69-87.

[6] G.K. Bakshi, M. Raka, A class of constacyclic codes over a finite field, Finite Fields Appl. 18 (2012) 362-377.

[7] B. Chen, H.Q. Dinh, H. Liu, Repeated-root constacyclic codes of length lp^s and their duals, Discrete Applied Math. 177 (2014) 60-70.

[8] Anuradha Sharma, Repeated-root constacyclic codes of length $l^t p^s$ and their duals codes, Cryptogr. Commun. 7 (2015) 229-255.

[9] B. Chen, H.Q. Dinh, H. Liu, Repeated-root constacyclic codes of length $2l^m p^s$, Finite Fields Appl. 33 (2015) 137-159.

[10] Y. Jia, S. Ling, C. Xing, On self-dual cyclic codes over finite fields, IEEE Trans. Inf. Theory 57 (2011) 2243-2251.

[11] X. Kai, S. Zhu, On cyclic self-dual codes, Appl. Algebra Engrg. Comm. Comput. 19 (2008) 509-525.

[12] H.Q. Dinh, On the lineare odering of some classess of negacyclic and cyclic codes and their distance distributions, Finite Fields Appl. 14 (2008) 22-40.

[13] H.Q. Dinh, Constacyclic codes of length p^s over $F_{p^m} + uF_{p^m}$, J. Algebra 324 (2010) 940-950.

[14] X. Kai, S. Zhu, On the distance of cyclic codes of length 2^e over Z_4, Discrete Math. 310 (2010) 12-20.

[15] A. Sharma, G.K. Bakshi, V.C. Dumir, M. Raka, Cyclotomic numbers and primitive idempotents in the ring $GF(q)[X]/\langle X^{p^n} - 1\rangle$, Finite Fields Appl. 10 (2004) 653-673.

[16] David M. Burton, Elementary Number Theory, Tata McGraw-Hill, 2006.

[17] Z. Wan, Lectures on Finite Fields and Galois Rings. Singapore: World Scientific Publishing, 2003.